


# The Eurasian Association on Inverse Problems (EAIP) Introducing The 2020 EAIP Award Winners



Jan Boman

*On Behalf of the EAIP Award Committee Members:*

*Alemdar Hasanov Hasanoglu, Bernd Hofmann,  
Vladimir G. Romanov, Gen Nakamura, Andreas Neubauer*



# Former EAIP Awardees

## ① The 2014 EAIP Award Winners:

Bernd Hofmann, Germany

Vladimir Vasin, Russian Federation



# Former EAIP Awardees

## ① The 2014 EAIP Award Winners:

Bernd Hofmann, Germany

Vladimir Vasin, Russian Federation

## ② 2016 EAIP Awardees:

Jan Boman, Sweden

Segrey I. Kabanikhin, Russian Federation



# Former EAIP Awardees

## ① The 2014 EAIP Award Winners:

Bernd Hofmann, Germany

Vladimir Vasin, Russian Federation

## ② 2016 EAIP Awardees:

Jan Boman, Sweden

Segrey I. Kabanikhin, Russian Federation

## ③ 2018 EAIP Awardees:

Vladimir G. Romanov, Russian Federation

Otmar Scherzer, Austria



## 2020 EAIP Award Recipients



Andreas Kirsch  
Germany



Roman Novikov  
France



Masahiro Yamamoto  
Japan



# Distinguished Contributions of Andreas Kirsch. I

- 1 Prof. Andreas Kirsch has enormous contributions to inverse scattering problem in many aspects. Among those the *linear sampling method* (LSM) and the *factorization method* are the most famous ones.



# Distinguished Contributions of Andreas Kirsch. I

- 1 Prof. Andreas Kirsch has enormous contributions to inverse scattering problem in many aspects. Among those the *linear sampling method* (LSM) and the *factorization method* are the most famous ones.
- 2 Together with Prof. David Colton he started the LSM. This method is the oldest and most developed non-iterative reconstruction method in inverse scattering theory to identify the shape of a scatterer, and has also been familiar in the engineering community.



# Distinguished Contributions of Andreas Kirsch. I

- 1 Prof. Andreas Kirsch has enormous contributions to inverse scattering problem in many aspects. Among those the *linear sampling method* (LSM) and the *factorization method* are the most famous ones.
- 2 Together with Prof. David Colton he started the LSM. This method is the oldest and most developed non-iterative reconstruction method in inverse scattering theory to identify the shape of a scatterer, and has also been familiar in the engineering community.
- 3 In order to completely validate this method mathematically, he developed more sophisticated method, the *factorization method*.





# Distinguished Contributions of Andreas Kirsch. I

- 1 Prof. Andreas Kirsch has enormous contributions to inverse scattering problem in many aspects. Among those the *linear sampling method* (LSM) and the *factorization method* are the most famous ones.
- 2 Together with Prof. David Colton he started the LSM. This method is the oldest and most developed non-iterative reconstruction method in inverse scattering theory to identify the shape of a scatterer, and has also been familiar in the engineering community.
- 3 In order to completely validate this method mathematically, he developed more sophisticated method, the *factorization method*.
- 4 This new method gave a strong impact to so many mathematicians and lead them to apply it to many equations in different setups and study its further generalization in many ways in the past 20 years.



# Distinguished Contributions of Andreas Kirsch. I

- ① Prof. Andreas Kirsch has enormous contributions to inverse scattering problem in many aspects. Among those the *linear sampling method* (LSM) and the *factorization method* are the most famous ones.
- ② Together with Prof. David Colton he started the LSM. This method is the oldest and most developed non-iterative reconstruction method in inverse scattering theory to identify the shape of a scatterer, and has also been familiar in the engineering community.
- ③ In order to completely validate this method mathematically, he developed more sophisticated method, the *factorization method*.
- ④ This new method gave a strong impact to so many mathematicians and lead them to apply it to many equations in different setups and study its further generalization in many ways in the past 20 years.
- ⑤ His book entitled *An Introduction to the Mathematical Theory of Inverse Problems*, (Springer 1996; 2nd edition 2011), became a standard work for introducing readers to the inverse problems.



## Distinguished Contributions of Andreas Kirsch. II

- 1 As a university professor and teacher at the Department of Mathematics of the Karlsruhe Institute of Technology, Prof. Andreas Kirsch inspired many young scientists for this field and for mathematics in general.



## Distinguished Contributions of Andreas Kirsch. II

- 1 As a university professor and teacher at the Department of Mathematics of the Karlsruhe Institute of Technology, Prof. Andreas Kirsch inspired many young scientists for this field and for mathematics in general.
- 2 He has a special reputation in the inverse problems community.



## Distinguished Contributions of Andreas Kirsch. II

- 1 As a university professor and teacher at the Department of Mathematics of the Karlsruhe Institute of Technology, Prof. Andreas Kirsch inspired many young scientists for this field and for mathematics in general.
- 2 He has a special reputation in the inverse problems community.
- 3 His strong affinity to international cooperation is especially to be emphasized.



## Distinguished Contributions of Andreas Kirsch. II

- ① As a university professor and teacher at the Department of Mathematics of the Karlsruhe Institute of Technology, Prof. Andreas Kirsch inspired many young scientists for this field and for mathematics in general.
- ② He has a special reputation in the inverse problems community.
- ③ His strong affinity to international cooperation is especially to be emphasized.
- ④ He coauthored a series of works with colleagues from different countries of Europe and Asia.



## Distinguished Contributions of Andreas Kirsch. II

- ① As a university professor and teacher at the Department of Mathematics of the Karlsruhe Institute of Technology, Prof. Andreas Kirsch inspired many young scientists for this field and for mathematics in general.
- ② He has a special reputation in the inverse problems community.
- ③ His strong affinity to international cooperation is especially to be emphasized.
- ④ He coauthored a series of works with colleagues from different countries of Europe and Asia.
- ⑤ Moreover, he provided a comprehensive service for the international inverse problems community in various fields of activity.



## Distinguished Contributions of Andreas Kirsch. II

- ① As a university professor and teacher at the Department of Mathematics of the Karlsruhe Institute of Technology, Prof. Andreas Kirsch inspired many young scientists for this field and for mathematics in general.
- ② He has a special reputation in the inverse problems community.
- ③ His strong affinity to international cooperation is especially to be emphasized.
- ④ He coauthored a series of works with colleagues from different countries of Europe and Asia.
- ⑤ Moreover, he provided a comprehensive service for the international inverse problems community in various fields of activity.
- ⑥ **The EAIP Award Committee has concluded that Prof. Andreas Kirsch deserves The Award of The Eurasian Association on Inverse Problems.**





# Distinguished Contributions of Roman Novikov. I

- ① In 1954 at the International Congress of Mathematicians held at Amsterdam, I.M.Gel'fand has formulated the following inverse problem for the Schrödinger equation  $-\Delta\psi + V(x) \cdot \psi = k^2\psi$ ,  $x \in \Omega \subset \mathbf{R}^3$ , where  $\Omega$  is a bounded domain.



# Distinguished Contributions of Roman Novikov. I

- ① In 1954 at the International Congress of Mathematicians held at Amsterdam, I.M.Gel'fand has formulated the following inverse problem for the Schrödinger equation  $-\Delta\psi + V(x) \cdot \psi = k^2\psi$ ,  $x \in \Omega \subset \mathbf{R}^3$ , where  $\Omega$  is a bounded domain.
- ② Can the potential  $V(x)$  be determined by the operator  $\mathcal{R}(k)$ , which is defined by Gelfand as the operator which *"establishes a relationship between the function  $\psi(x)$  and its normal derivative  $\partial\psi/\partial\nu$  on the surface  $\partial\Omega$ "*, and if yes, how?



# Distinguished Contributions of Roman Novikov. I

- ① In 1954 at the International Congress of Mathematicians held at Amsterdam, I.M.Gel'fand has formulated the following inverse problem for the Schrödinger equation  $-\Delta\psi + V(x) \cdot \psi = k^2\psi$ ,  $x \in \Omega \subset \mathbf{R}^3$ , where  $\Omega$  is a bounded domain.
- ② Can the potential  $V(x)$  be determined by the operator  $\mathcal{R}(k)$ , which is defined by Gelfand as the operator which "*establishes a relationship between the function  $\psi(x)$  and its normal derivative  $\partial\psi/\partial\nu$  on the surface  $\partial\Omega$* ", and if yes, how?
- ③ See [I.M.Gel'fand, Some Problems of Functional Analysis and Algebra, *Proceedings of the International Congress of Mathematicians*, Amsterdam, Volume I, pp. 253-276 (1954)]



# Distinguished Contributions of Roman Novikov. I

- ① In 1954 at the International Congress of Mathematicians held at Amsterdam, I.M.Gel'fand has formulated the following inverse problem for the Schrödinger equation  $-\Delta\psi + V(x) \cdot \psi = k^2\psi$ ,  $x \in \Omega \subset \mathbf{R}^3$ , where  $\Omega$  is a bounded domain.
- ② Can the potential  $V(x)$  be determined by the operator  $\mathcal{R}(k)$ , which is defined by Gelfand as the operator which "*establishes a relationship between the function  $\psi(x)$  and its normal derivative  $\partial\psi/\partial\nu$  on the surface  $\partial\Omega$* ", and if yes, how?
- ③ See [I.M.Gel'fand, Some Problems of Functional Analysis and Algebra, *Proceedings of the International Congress of Mathematicians*, Amsterdam, Volume I, pp. 253-276 (1954)]
- ④ This work opened a new research field in inverse problems, namely, determination of an unknown coefficient from the Dirichlet-to-Neumann operator.



# Distinguished Contributions of Roman Novikov. I

- ① In 1954 at the International Congress of Mathematicians held at Amsterdam, I.M.Gel'fand has formulated the following inverse problem for the Schrödinger equation  $-\Delta\psi + V(x) \cdot \psi = k^2\psi$ ,  $x \in \Omega \subset \mathbf{R}^3$ , where  $\Omega$  is a bounded domain.
- ② Can the potential  $V(x)$  be determined by the operator  $\mathcal{R}(k)$ , which is defined by Gelfand as the operator which "establishes a relationship between the function  $\psi(x)$  and its normal derivative  $\partial\psi/\partial\nu$  on the surface  $\partial\Omega$ ", and if yes, how?
- ③ See [I.M.Gel'fand, Some Problems of Functional Analysis and Algebra, *Proceedings of the International Congress of Mathematicians*, Amsterdam, Volume I, pp. 253-276 (1954)]
- ④ This work opened a new research field in inverse problems, namely, determination of an unknown coefficient from the Dirichlet-to-Neumann operator.
- ⑤ After 33 years, the Gelfand's problem in its original formulation has been solved by R.G. Novikov in [R.G. Novikov, G.M. Khenkin, The  $\bar{\partial}$ -equation in the multidimensional inverse scattering problem, *Uspehki Mat. Nauk*, **42**:3 (1987), 93-152],



# Distinguished Contributions of Roman Novikov. I

- ① In 1954 at the International Congress of Mathematicians held at Amsterdam, I.M.Gel'fand has formulated the following inverse problem for the Schrödinger equation  $-\Delta\psi + V(x) \cdot \psi = k^2\psi$ ,  $x \in \Omega \subset \mathbf{R}^3$ , where  $\Omega$  is a bounded domain.
- ② Can the potential  $V(x)$  be determined by the operator  $\mathcal{R}(k)$ , which is defined by Gelfand as the operator which "establishes a relationship between the function  $\psi(x)$  and its normal derivative  $\partial\psi/\partial\nu$  on the surface  $\partial\Omega$ ", and if yes, how?
- ③ See [I.M.Gel'fand, Some Problems of Functional Analysis and Algebra, *Proceedings of the International Congress of Mathematicians*, Amsterdam, Volume I, pp. 253-276 (1954)]
- ④ This work opened a new research field in inverse problems, namely, determination of an unknown coefficient from the Dirichlet-to-Neumann operator.
- ⑤ After 33 years, the Gelfand's problem in its original formulation has been solved by R.G. Novikov in [R.G. Novikov, G.M. Khenkin, The  $\bar{\partial}$ -equation in the multidimensional inverse scattering problem, *Uspehki Mat. Nauk*, **42**:3 (1987), 93-152],
- ⑥ when he was a first year PhD student at the Moscow State University, beeing only 22 years old.



## Distinguished Contributions of Roman Novikov. II

- 1 In today's terminology, Prof. Roman Novikov *has first solved one of the fundamental problems of inverse problems theory*, namely the problem of invertibility of the Dirichlet-to-Neumann operator corresponding to inverse coefficient problem for the Schrödinger equation.



## Distinguished Contributions of Roman Novikov. II

- 1 In today's terminology, Prof. Roman Novikov *has first solved one of the fundamental problems of inverse problems theory*, namely the problem of invertibility of the Dirichlet-to-Neumann operator corresponding to inverse coefficient problem for the Schrödinger equation.
- 2 The "*Novikov inversion formula*", proposed exactly 20 years ago has become now a classic.





## Distinguished Contributions of Roman Novikov. II

- 1 In today's terminology, Prof. Roman Novikov *has first solved one of the fundamental problems of inverse problems theory*, namely the problem of invertibility of the Dirichlet-to-Neumann operator corresponding to inverse coefficient problem for the Schrödinger equation.
- 2 The "*Novikov inversion formula*", proposed exactly 20 years ago has become now a classic.
- 3 As a university professor and Project Director at the Center for Applied Mathematics of Ecole Polytechnique, he inspired many young scientists from Europe and Asia.



## Distinguished Contributions of Roman Novikov. II

- ① In today's terminology, Prof. Roman Novikov *has first solved one of the fundamental problems of inverse problems theory*, namely the problem of invertibility of the Dirichlet-to-Neumann operator corresponding to inverse coefficient problem for the Schrödinger equation.
- ② The "*Novikov inversion formula*", proposed exactly 20 years ago has become now a classic.
- ③ As a university professor and Project Director at the Center for Applied Mathematics of Ecole Polytechnique, he inspired many young scientists from Europe and Asia.
- ④ Prof. Roman Novikov has distinguished students with high scientific level and impact. Some of them were awarded by the EAIP Young Scientist Award.



## Distinguished Contributions of Roman Novikov. II

- ① In today's terminology, Prof. Roman Novikov *has first solved one of the fundamental problems of inverse problems theory*, namely the problem of invertibility of the Dirichlet-to-Neumann operator corresponding to inverse coefficient problem for the Schrödinger equation.
- ② The "*Novikov inversion formula*", proposed exactly 20 years ago has become now a classic.
- ③ As a university professor and Project Director at the Center for Applied Mathematics of Ecole Polytechnique, he inspired many young scientists from Europe and Asia.
- ④ Prof. Roman Novikov has distinguished students with high scientific level and impact. Some of them were awarded by the EAIP Young Scientist Award.
- ⑤ For many years, he has made important contributions to international meetings on inverse problems. He is one of the prominent members of the inverse problems community.



## Distinguished Contributions of Roman Novikov. II

- ① In today's terminology, Prof. Roman Novikov *has first solved one of the fundamental problems of inverse problems theory*, namely the problem of invertibility of the Dirichlet-to-Neumann operator corresponding to inverse coefficient problem for the Schrödinger equation.
- ② The "*Novikov inversion formula*", proposed exactly 20 years ago has become now a classic.
- ③ As a university professor and Project Director at the Center for Applied Mathematics of Ecole Polytechnique, he inspired many young scientists from Europe and Asia.
- ④ Prof. Roman Novikov has distinguished students with high scientific level and impact. Some of them were awarded by the EAIP Young Scientist Award.
- ⑤ For many years, he has made important contributions to international meetings on inverse problems. He is one of the prominent members of the inverse problems community.
- ⑥ The EAIP Award Committee has concluded that Prof. Roman Novikov deserves **The Award of The Eurasian Association on Inverse Problems.**



# Distinguished Contributions of Masahiro Yamamoto. I

- 1 Prof. Masahiro Yamamoto has contributed a lot to many areas of inverse problems, in the past 30 years.



# Distinguished Contributions of Masahiro Yamamoto. I

- 1 Prof. Masahiro Yamamoto has contributed a lot to many areas of inverse problems, in the past 30 years.
- 2 In different kinds of inverse problems, one can see the original approach inherent in Professor Yamamoto.



# Distinguished Contributions of Masahiro Yamamoto. I

- 1 Prof. Masahiro Yamamoto has contributed a lot to many areas of inverse problems, in the past 30 years.
- 2 In different kinds of inverse problems, one can see the original approach inherent in Professor Yamamoto.
- 3 In the most well-known data base mathscinet in mathematics, the total number of his publications is over 325 and the total number of citations is over 4400, which is really incredible.



# Distinguished Contributions of Masahiro Yamamoto. I

- 1 Prof. Masahiro Yamamoto has contributed a lot to many areas of inverse problems, in the past 30 years.
- 2 In different kinds of inverse problems, one can see the original approach inherent in Professor Yamamoto.
- 3 In the most well-known data base mathscinet in mathematics, the total number of his publications is over 325 and the total number of citations is over 4400, which is really incredible.
- 4 Among them, Prof. Yamamoto achieved significant results, especially with regard to the following three classes of problems.





# Distinguished Contributions of Masahiro Yamamoto. I

- 1 Prof. Masahiro Yamamoto has contributed a lot to many areas of inverse problems, in the past 30 years.
- 2 In different kinds of inverse problems, one can see the original approach inherent in Professor Yamamoto.
- 3 In the most well-known data base mathscinet in mathematics, the total number of his publications is over 325 and the total number of citations is over 4400, which is really incredible.
- 4 Among them, Prof. Yamamoto achieved significant results, especially with regard to the following three classes of problems.
- 5 Application of Carleman estimates showing the uniqueness and stability for the inverse boundary value problems.



# Distinguished Contributions of Masahiro Yamamoto. I

- ① Prof. Masahiro Yamamoto has contributed a lot to many areas of inverse problems, in the past 30 years.
- ② In different kinds of inverse problems, one can see the original approach inherent in Professor Yamamoto.
- ③ In the most well-known data base mathscinet in mathematics, the total number of his publications is over 325 and the total number of citations is over 4400, which is really incredible.
- ④ Among them, Prof. Yamamoto achieved significant results, especially with regard to the following three classes of problems.
- ⑤ Application of Carleman estimates showing the uniqueness and stability for the inverse boundary value problems.
- ⑥ Triggered and developed the study of inverse problems for diffusion equation with time fractional derivatives.



# Distinguished Contributions of Masahiro Yamamoto. I

- ① Prof. Masahiro Yamamoto has contributed a lot to many areas of inverse problems, in the past 30 years.
- ② In different kinds of inverse problems, one can see the original approach inherent in Professor Yamamoto.
- ③ In the most well-known data base mathscinet in mathematics, the total number of his publications is over 325 and the total number of citations is over 4400, which is really incredible.
- ④ Among them, Prof. Yamamoto achieved significant results, especially with regard to the following three classes of problems.
- ⑤ Application of Carleman estimates showing the uniqueness and stability for the inverse boundary value problems.
- ⑥ Triggered and developed the study of inverse problems for diffusion equation with time fractional derivatives.
- ⑦ Several successful studies on industrial inverse problems.



## Distinguished Contributions of Masahiro Yamamoto. II

- 1 As a university professor and teacher at the Graduate School of Mathematical Sciences of The University of Tokyo he motivated many young scientists to study applied mathematics and in particular inverse problems.



## Distinguished Contributions of Masahiro Yamamoto. II

- 1 As a university professor and teacher at the Graduate School of Mathematical Sciences of The University of Tokyo he motivated many young scientists to study applied mathematics and in particular inverse problems.
- 2 Prof. Masahiro Yamamoto is one of the most traveling colleagues in the international inverse problems community. In this context and as an associate and guest editor of a couple of renowned international journals, he did much for the cooperation between Asia and Europe.



## Distinguished Contributions of Masahiro Yamamoto. II

- 1 As a university professor and teacher at the Graduate School of Mathematical Sciences of The University of Tokyo he motivated many young scientists to study applied mathematics and in particular inverse problems.
- 2 Prof. Masahiro Yamamoto is one of the most traveling colleagues in the international inverse problems community. In this context and as an associate and guest editor of a couple of renowned international journals, he did much for the cooperation between Asia and Europe.
- 3 He also succeeded in establishing a large group for this study in China by educating many European and Chinese Ph.D. students at the University of Tokyo.



## Distinguished Contributions of Masahiro Yamamoto. II

- 1 As a university professor and teacher at the Graduate School of Mathematical Sciences of The University of Tokyo he motivated many young scientists to study applied mathematics and in particular inverse problems.
- 2 Prof. Masahiro Yamamoto is one of the most traveling colleagues in the international inverse problems community. In this context and as an associate and guest editor of a couple of renowned international journals, he did much for the cooperation between Asia and Europe.
- 3 He also succeeded in establishing a large group for this study in China by educating many European and Chinese Ph.D. students at the University of Tokyo.
- 4 The outcomes of these achievements are really nice giving good evidences how mathematics can be effective in solving some practical inverse problems



## Distinguished Contributions of Masahiro Yamamoto. II

- ① As a university professor and teacher at the Graduate School of Mathematical Sciences of The University of Tokyo he motivated many young scientists to study applied mathematics and in particular inverse problems.
- ② Prof. Masahiro Yamamoto is one of the most traveling colleagues in the international inverse problems community. In this context and as an associate and guest editor of a couple of renowned international journals, he did much for the cooperation between Asia and Europe.
- ③ He also succeeded in establishing a large group for this study in China by educating many European and Chinese Ph.D. students at the University of Tokyo.
- ④ The outcomes of these achievements are really nice giving good evidences how mathematics can be effective in solving some practical inverse problems
- ⑤ **The EAIP Award Committee has concluded that Prof. Masahiro Yamamoto deserves The Award of The Eurasian Association on Inverse Problems.**

